

# Detection of B-mode Polarization at 150 GHz and Degree Angular Scales by BICEP2 and Keck Array

# Modern cosmology in a nutshell:



Edwin Hubble

1) The universe is expanding.  
(Hubble, 1920s)

2) It was once hot and dense, like the inside of the Sun.

(Alpher, Gamow, Herman, 1940s)

3) You can still see the glow!  
*The Cosmic Microwave Background*  
(Penzias & Wilson, 1964)



Bob Wilson & Arno Penzias  
1978 Nobel Prize

⇒ acceptance of the “**HOT BIG BANG**”



**INFLATION**

**fraction  
of a second**

**CMB  
last scattering**


**379,000  
years**

**first  
stars**


**~200 million  
years**

**present  
day**


**13.7 billion  
years**



# History of the Universe




# History of the Universe







# History of the Universe



# History of the Universe



# History of the Universe





# CMB Temperature Measurements / Inflation

CMB temperature anisotropy now measured over full range of angular scales.



Consistent with  $\Lambda$ CDM paradigm(?) and constrains its parameters to sub percent accuracy.

Inflation “invented” in 1980s to explain facts about the Universe which were known or suspected.

Makes additional prediction of a background of gravitational waves (aka tensor modes) – which will imprint a specific CMB polarization pattern...

→ so-called “smoking gun”

→ amplitude tells us the energy scale at which inflation occurred



# Why Inflation?

Solves the horizon problem:  
Why is the CMB nearly uniform?  
How do apparently causally disconnected regions of space get set to the same temperature?



A volume much larger than our entire observable universe today was once a causally connected sub atomic spec.

Solves the flatness problem:  
Why is the net spatial curvature close to zero?



Any initial spatial curvature is diluted away to undetectability by the hyper expansion.

Explains the initial perturbation spectrum: Why was it close to flat power law?




Equal amount of perturbations are injected at each step in the exponential expansion.

Solves the monopole problem:  
Why do we not observe magnetic monopoles in the Universe today?



Monopoles are diluted away to undetectability.


**CMB polarization:  
arises at last scattering  
from local radiation quadrupole**






# CMB polarization


Density Wave




E-Mode Polarization Pattern




Gravitational Wave



B-Mode Polarization Pattern



# The State of B-mode Measurements last March



In simple inflationary gravitational wave models the

**tensor-to-scalar ratio  $r$**

is the only parameter to the B-mode spectrum.

Before BICEP2: only upper limits from searches for Inflationary B-modes

BICEP1 limits translated to:

**$r < 0.7$  (95% CL)**

At high multipoles lensing B-mode dominant.

SPT x-corr: lower limits on lensing B-mode from cross correlation using the CIB





UNIVERSITY OF  
TORONTO





# The BICEP2/Keck Postdocs



Colin Bischoff



Jeff Filippini




Martin Lueker



Walt Ogburn



Abigail Viereggen



Immanuel Buder



Stefan Fliescher



Roger O'Brient



Angiola Orlando




Zak Staniszewski

# The BICEP2/Keck Graduate Students



Randol Aikin




Justus Brevik



Chris Sheehy



Grant Teply




Chin Lin Wong



Kirit Karkare



Jon Kaufman



Sarah Kernasovskiy



Jamie Tolan

## Winterovers

### BICEP2

Steffen Richter

2010

### Keck



Steffen Richter

2011



Robert Schwarz



Steffen Richter

2012



Robert Schwarz

2013



Robert Schwarz

2014



Robert Schwarz

2015



Robert Schwarz

# South Pole CMB telescopes



NSF's South Pole Station:  
A popular place with CMB Experimentalists!

Super dry atmosphere and 24h coverage of low foreground sky.  
Also power, LHe,  $\text{LN}_2$ , 200 GB/day, 3 square meals, and bingo night...





**BICEP1**  
(2006 - 8)

**BICEP2**  
(2010 - 12)




**Keck Array**  
(2011 -)

**BICEP3**  
(2014 -)





Telescope and Mount



Focal Plane



Beams on Sky



-5 0 5  
Longitude (degrees)


-5 0 5  
Longitude (degrees)

-5 0 5  
Longitude (degrees)

-10 -5 0 5 10  
Longitude (degrees)




# BICEP2/Keck Experimental Concept

- Small aperture
- Wide field of view
- Cold refractor





# Mass-produced superconducting detectors



Slot antennas





# Observational Strategy



Go deep in a region of sky where galactic foregrounds are low


Observe at frequencies where the CMB is brightest with respect to:

Synchrotron emission (from high energy electrons)  
- falls with increasing freq

Thermal dust emission – rises with increasing freq

Foreground contamination of the B-mode power in clean regions previously projected to be equivalent to  $r \leq \sim 0.01$ .






Clem Pryke for The Bicep2 Collaboration

Time 50 mins

Time 00 Time 1




- Cover the whole field in 60 such scansets then start over at new boresight rotation
- Scanning modulates the CMB signal to freqs  $< 4$  Hz

# Raw Data - Worse Weather

Time 50 mins




Run 20120622D01\_dk293, scan block 1; Rx: 0





➤ Scanning over lumpy atmosphere  
→ “clouds”

➤ Pair difference still clean  
→ atmosphere is unpolarized


# BICEP2 3-year Data Set







# Total Polarization




E-mode dominated pattern – no obvious curl component

# B-mode Contribution

BICEP2 B-mode signal

Scale:  $1.7 \mu K$




Apply purification operation to Q/U maps which leaves only B-modes (given all timestream filterings etc.)

# B-mode Contribution


BICEP2 B-mode signal

Scale:  $0.3 \mu K$





Stretch scale by factor 6 – see “swirly” B-mode

# BICEP2 B-mode Power Spectrum





# Pre-Planck Polarized Dust Foreground Projections




The BICEP2 region was chosen on the basis of extremely low *unpolarized* dust power.

Used various models of polarized dust emission to estimate dust power.

Result: All auto spectra were well below observed signal level. (and cross spectra consistent with zero.)

But considerable uncertainty in these models...


# Fitting with Dust Projections Subtracted...



Probability that each of these models reflected reality was hard to assess.


DDM1 used all publicly available information from Planck. Polarization fraction here assumed  $p = 5\%$ .  $p \sim 13\%$  would explain the full excess under this model.

Adjust likelihood curve by subtracting the dust projection auto and cross spectra from our bandpowers:



# Conclusions circa March 17<sup>th</sup> 2014

BICEP2 data and upper limits from other experiments:



Most sensitive polarization maps ever made!

Power spectra perfectly consistent with lensed- $\Lambda$ CDM except:  
 $5.2\sigma$  excess in the B-mode spectrum at low multipoles!

Extensive studies and jackknife tests strongly argued against systematics as the origin

Data fit well to LCDM+ $r=0.2$  expectation

Foregrounds did not appear to be a large fraction of the signal...



# Storm of Media Attention

12.00 THE NOTICES NEWS

**USA TODAY**  
03.18.14  
A GANNETT COMPANY

**NCAA TOURNAMENT**  
WHO HAS BEST "DANCE CARDS"  
A look at matchups, players and teams to watch, 5C

**UConn tops women's tourney**  
ANALYSIS, BRACKET, 4C

## Putin, U.S. up ante after vote

Sanctions imposed, Ukraine, Russia ready troops as Duma considers Crimean annexation

**'Always hope' missing jet's passengers alive**  
As search expands to two helicopters, Malaysian officials won't rule out possibility that plane is intact. **3A**

**GM issues three new recalls**  
New recalls involve all 2014 Buick, Buick system plug, involve more than 15 million vehicles. **1B**

**Homework load unchanged**  
Despite parents' concerns about more work, study finds burden has barely changed over 30 years. **3A**

**How earnings earner big, bad wings**  
To play "The Wolf of Wall Street" as a CEO, DiCaprio's critics won't let him do it. **1D**

**South Pole view**  
The first photo taken from the South Pole, where the majority of our energy is produced, is a new look at the continent. **3D**

**U.S. SNAPSHOTS**  
A look at the day's news, from the White House to the streets. **1C**

**WAVES COULD BE BIG BANG'S SMOKING GUN**  
Scientists of the South Pole used special telescopes to detect primordial gravitational waves—ripples in the fabric of space and time—which hold clues to the nature of the universe. The ripples have never been seen directly until now. **Story 5A**

**Pen energy**  
Scientists are trying to prove that gravitational waves exist by detecting the tiny ripples in space and time that are predicted to exist. **Story 5A**

**Gravitational waves**  
Scientists have been searching for gravitational waves, ripples in the fabric of space and time, for decades. Now, they may have found them. **Story 5A**

**A THEORY**  
In 1915, Albert Einstein's theory of general relativity predicted that gravity would warp the fabric of space and time. Now, scientists are trying to prove it. **Story 5A**

**States engage in shadowy deals as death penalty drugs dwindle**  
Prisoners have drug, use shady pharmacies, try untested methods. **Story 5A**

**Prisoners have drug, use shady pharmacies, try untested methods**  
Prisoners have drug, use shady pharmacies, try untested methods. **Story 5A**

**Prisoners have drug, use shady pharmacies, try untested methods**  
Prisoners have drug, use shady pharmacies, try untested methods. **Story 5A**

**Prisoners have drug, use shady pharmacies, try untested methods**  
Prisoners have drug, use shady pharmacies, try untested methods. **Story 5A**

**Prisoners have drug, use shady pharmacies, try untested methods**  
Prisoners have drug, use shady pharmacies, try untested methods. **Story 5A**

**Prisoners have drug, use shady pharmacies, try untested methods**  
Prisoners have drug, use shady pharmacies, try untested methods. **Story 5A**

**Prisoners have drug, use shady pharmacies, try untested methods**  
Prisoners have drug, use shady pharmacies, try untested methods. **Story 5A**

**Prisoners have drug, use shady pharmacies, try untested methods**  
Prisoners have drug, use shady pharmacies, try untested methods. **Story 5A**

"All the News That's Fit to Print"

**The New York Times**

VOL. CLXXIII, No. 56,444  
TUESDAY, MARCH 18, 2014

New England Edition  
\$2.50



**PUTIN RECOGNIZES CRIMEA SECESSION, DEFEYING THE WEST**  
Decree Increases Fears of Annexation by Russia, Despite More Sanctions

BY STEVEN LEVITSKY AND PETER BAKER  
Moscow — President Vladimir Putin's decree recognizing Crimea's secession from Ukraine, a move that defies the West, has triggered a new round of international sanctions against Russia. The move also signals a major shift in Russian foreign policy, as the country's leader moves to assert his authority over the country's actions.

**Space Ripples Reveal Big Bang's Smoking Gun**  
Scientists have found evidence of gravitational waves, ripples in the fabric of space and time, which hold clues to the nature of the universe. The ripples have never been seen directly until now.

**Lost Jet's Path Seen as Altered**  
A new study suggests that the path of the missing Malaysia Airlines flight may have been altered by a strong jet stream, which could have led to the plane's crash.

**States engage in shadowy deals as death penalty drugs dwindle**  
Prisoners have drug, use shady pharmacies, try untested methods.

**Prisoners have drug, use shady pharmacies, try untested methods**  
Prisoners have drug, use shady pharmacies, try untested methods.

**Prisoners have drug, use shady pharmacies, try untested methods**  
Prisoners have drug, use shady pharmacies, try untested methods.

**Prisoners have drug, use shady pharmacies, try untested methods**  
Prisoners have drug, use shady pharmacies, try untested methods.

**Prisoners have drug, use shady pharmacies, try untested methods**  
Prisoners have drug, use shady pharmacies, try untested methods.

**Prisoners have drug, use shady pharmacies, try untested methods**  
Prisoners have drug, use shady pharmacies, try untested methods.

**Prisoners have drug, use shady pharmacies, try untested methods**  
Prisoners have drug, use shady pharmacies, try untested methods.

**Prisoners have drug, use shady pharmacies, try untested methods**  
Prisoners have drug, use shady pharmacies, try untested methods.

**Prisoners have drug, use shady pharmacies, try untested methods**  
Prisoners have drug, use shady pharmacies, try untested methods.

**Prisoners have drug, use shady pharmacies, try untested methods**  
Prisoners have drug, use shady pharmacies, try untested methods.

**Prisoners have drug, use shady pharmacies, try untested methods**  
Prisoners have drug, use shady pharmacies, try untested methods.



Image courtesy of NASA. The satellite dish is part of the Gravity Probe B mission, which is designed to test Einstein's theory of general relativity.

## 宇宙急速膨張の証拠、検出される

Telescope captures view of gravitational waves

Russ. Cosmos 2014年3月18日 09時 46分 507 (2014-03-18)

宇宙が急速に膨張しているという証拠(インフレーション)として、重力波の検出が期待されている。重力波は、宇宙の膨張に伴って発生する波であり、重力波の検出は、宇宙の膨張の速度を測定するのに役立つ。

重力波は、宇宙の膨張に伴って発生する波であり、重力波の検出は、宇宙の膨張の速度を測定するのに役立つ。

重力波は、宇宙の膨張に伴って発生する波であり、重力波の検出は、宇宙の膨張の速度を測定するのに役立つ。

重力波は、宇宙の膨張に伴って発生する波であり、重力波の検出は、宇宙の膨張の速度を測定するのに役立つ。

重力波は、宇宙の膨張に伴って発生する波であり、重力波の検出は、宇宙の膨張の速度を測定するのに役立つ。

重力波は、宇宙の膨張に伴って発生する波であり、重力波の検出は、宇宙の膨張の速度を測定するのに役立つ。

重力波は、宇宙の膨張に伴って発生する波であり、重力波の検出は、宇宙の膨張の速度を測定するのに役立つ。

重力波は、宇宙の膨張に伴って発生する波であり、重力波の検出は、宇宙の膨張の速度を測定するのに役立つ。

重力波は、宇宙の膨張に伴って発生する波であり、重力波の検出は、宇宙の膨張の速度を測定するのに役立つ。

重力波は、宇宙の膨張に伴って発生する波であり、重力波の検出は、宇宙の膨張の速度を測定するのに役立つ。

重力波は、宇宙の膨張に伴って発生する波であり、重力波の検出は、宇宙の膨張の速度を測定するのに役立つ。

重力波は、宇宙の膨張に伴って発生する波であり、重力波の検出は、宇宙の膨張の速度を測定するのに役立つ。

重力波は、宇宙の膨張に伴って発生する波であり、重力波の検出は、宇宙の膨張の速度を測定するのに役立つ。

重力波は、宇宙の膨張に伴って発生する波であり、重力波の検出は、宇宙の膨張の速度を測定するのに役立つ。

重力波は、宇宙の膨張に伴って発生する波であり、重力波の検出は、宇宙の膨張の速度を測定するのに役立つ。

重力波は、宇宙の膨張に伴って発生する波であり、重力波の検出は、宇宙の膨張の速度を測定するのに役立つ。

重力波は、宇宙の膨張に伴って発生する波であり、重力波の検出は、宇宙の膨張の速度を測定するのに役立つ。

重力波は、宇宙の膨張に伴って発生する波であり、重力波の検出は、宇宙の膨張の速度を測定するのに役立つ。

重力波は、宇宙の膨張に伴って発生する波であり、重力波の検出は、宇宙の膨張の速度を測定するのに役立つ。

重力波は、宇宙の膨張に伴って発生する波であり、重力波の検出は、宇宙の膨張の速度を測定するのに役立つ。

重力波は、宇宙の膨張に伴って発生する波であり、重力波の検出は、宇宙の膨張の速度を測定するのに役立つ。

重力波は、宇宙の膨張に伴って発生する波であり、重力波の検出は、宇宙の膨張の速度を測定するのに役立つ。

重力波は、宇宙の膨張に伴って発生する波であり、重力波の検出は、宇宙の膨張の速度を測定するのに役立つ。

重力波は、宇宙の膨張に伴って発生する波であり、重力波の検出は、宇宙の膨張の速度を測定するのに役立つ。

重力波は、宇宙の膨張に伴って発生する波であり、重力波の検出は、宇宙の膨張の速度を測定するのに役立つ。

重力波は、宇宙の膨張に伴って発生する波であり、重力波の検出は、宇宙の膨張の速度を測定するのに役立つ。

重力波は、宇宙の膨張に伴って発生する波であり、重力波の検出は、宇宙の膨張の速度を測定するのに役立つ。

重力波は、宇宙の膨張に伴って発生する波であり、重力波の検出は、宇宙の膨張の速度を測定するのに役立つ。

重力波は、宇宙の膨張に伴って発生する波であり、重力波の検出は、宇宙の膨張の速度を測定するのに役立つ。

重力波は、宇宙の膨張に伴って発生する波であり、重力波の検出は、宇宙の膨張の速度を測定するのに役立つ。

重力波は、宇宙の膨張に伴って発生する波であり、重力波の検出は、宇宙の膨張の速度を測定するのに役立つ。

重力波は、宇宙の膨張に伴って発生する波であり、重力波の検出は、宇宙の膨張の速度を測定するのに役立つ。

重力波は、宇宙の膨張に伴って発生する波であり、重力波の検出は、宇宙の膨張の速度を測定するのに役立つ。

重力波は、宇宙の膨張に伴って発生する波であり、重力波の検出は、宇宙の膨張の速度を測定するのに役立つ。

重力波は、宇宙の膨張に伴って発生する波であり、重力波の検出は、宇宙の膨張の速度を測定するのに役立つ。

112

**PHYSICAL REVIEW LETTERS**

Library or Other Institution Use Prohibited Until 2017

Articles published week ending 20 JUNE 2014

PHYS. REV. LETT. 112 (24 JUNE 2014) 240001-240010 (24 June 2014) (248 total pages)

24

Published by American Physical Society.

Volume 112, Number 24


Actually not a lot of fun...



# Developments last year

- Intense media and science community interest...
- Many early instrumental queries – faded away – everybody now seems to trust our measurements.
- Concerns about synchrotron – also faded away.
- But persistent concerns about dust...
  - Mostly based on online pdf's of Planck talks
- In September we finally got some solid information from Planck about the actual level of polarized dust emission in the BICEP2 field (arxiv:1409.5738). Much higher than any of the projections...


# Results from Joint analysis of BICEP2/Keck and Planck data



- In summer 2014 BICEP2/Keck and Planck collaborations signed MOU to do a joint analysis of their data
- Data exchanged in late July
- Today reporting on results of this analysis as presented in paper [arxiv:1502.00612](https://arxiv.org/abs/1502.00612) (now published by PRL)


← BICEP2/Keck/Planck  
meeting at University  
of Minnesota 5 Nov  
2014

# B2 150 GHz T/Q/U maps of small sky patch





# B2+Keck 150 GHz T/Q/U maps of small sky patch




57 nK deg (3.4  $\mu\text{K}$  arcmin) when adding 2012/13 Keck data -  
by far the deepest maps ever made - but apodized and  
filtered...

# Planck

WMAP

Planck

COBE




graphic: J. Gudmundsson

- Planck is the third space mission to observe the CMB: An ESA-led mission Launched 14 May 2009, mission completed Oct 2013
- Full sky maps produced in seven polarization-sensitive bands centered at 30,44,70,(100,143,217),353 GHz (to be) released in 2015. Also intensity maps at 545 and 857 GHz.



# Planck full sky maps at 9 frequencies





Full sky coverage and 9 frequencies - but not as deep as BICEP2/Keck in any given region of the sky



# Planck 353 GHz full sky maps in polarization


- 353 GHz polarized maps are dominated by Galactic dust emission

For comparison, Planck 70 GHz is close to the minimum of Galactic foreground emission




-20  $\mu\text{K}$  20

-20  $\mu\text{K}$  20




-100.0  $\mu\text{K}_{\text{CMB}}$  100.0  $\mu\text{K}_{\text{CMB}}$



-100.0  $\mu\text{K}_{\text{CMB}}$  100.0  $\mu\text{K}_{\text{CMB}}$

# Compare BK 150 GHz (left) with Planck 353 GHz (right)




Dominated by LCDM E-modes

Q/U maps x10 color stretch

Dominated by noise&dust

The Real Data

# Compare BK 150 GHz (left) with Planck 353 GHz (right)




A Noise Simulation



# What are the expectations for dust?

- In the BK patch Planck's signal-to-noise on dust is limited even at 353GHz.
- However a series of Planck papers have investigated the spatial and frequency spectra of dust over the intermediate and high latitude sky:


Fig 2 of arxiv:1409.5738



Dust BB spatial power spectra follow  $\ell^{-0.42}$  power law when averaging over large sky regions

- No evidence of deviation from this behavior for small sky patches although s/n low

Fig 6 of arxiv:1409.5738





Spectral energy distribution of polarized dust emission follows modified blackbody model with  $T=19.6\text{K}$  and  $\beta_d=1.59$

- Seems to be remarkably uniform over the high latitude sky

→ Good news for component separation

# Cross Correlation of BK and Planck 353 GHz




- Correlation of 150 GHz and 353 GHz B-modes is detected with high signal-to-noise.
- Scaling the cross-frequency spectrum by the expected brightness ratio (x25) of dust (right y-axis) indicates that dust contribution is comparable in magnitude to BICEP2/Keck excess over LCDM.
  - Shape looks consistent with  $\ell^{-0.42}$  power law expectation


# Multi-component multi-spectral likelihood analysis

- Define “fiducial analysis” to use single- and cross-frequency spectra between BK 150 GHz and Planck 217&353 GHz channels
  - (Detail: for Planck single-frequency use detector set split cross spectrum)
- As addition to basic LCDM lensing signal include gravity wave signal (with amp  $r$ ) and dust signal with amplitude  $A_d$  (specified at  $\ell = 80$  and 353 GHz)
  - For dust SED use modified blackbody model and marginalize over range  $\beta_d = 1.59 \pm 0.11$
- Use 5 lowest BB bandpowers only ( $20 < \ell < 200$ )


# Multi-component multi-spectral likelihood analysis



$r$  constraint consistent with zero (For BK+P  $L_0/L_{\text{peak}}$  ratio is 0.4 which happens 8% of the time in a dust only model.)




Dust is detected with  $5.1 \sigma$  significance




As expected dust and  $r$  are partially degenerate - reducing dust means more of the  $150 \times 150$  signal needs to be  $r$









# Best fit multi-frequency model



Model:




- The maximum likelihood model has acceptable  $\chi^2$  (with the biggest contribution coming from P353xP353.)
- The BKxBK and BKxP353 spectra are both very well fit by the model.



These plots show data as “naked points” versus center value and spread of best fit model to emphasize that uncertainty varies with the model (due to sample variance)

# Comparison of signal and noise levels




- The BICEP2/Keck noise is much lower than the Planck noise in the small sky patch observed
- However dust is much brighter at 353 GHz and Planck detects it
- The noise in the cross spectra is the geometric mean and a fairly tight constraint on dust amplitude is set

# Current Conclusions

- Last March BICEP2 reported detection of B-mode polarization in the CMB at 150GHz well in excess of the standard model expectation
  - This signal is confirmed by new data from the successor experiment Keck Array
- Last summer Planck released new information on the polarized emission from galactic dust which showed this might be due to dust emission.
- We have now done a joint analysis with Planck - The fundamental conclusion is that dust is detected at high significance, and  $r < 0.12$  at 95% confidence.
  - Multi-component likelihood gives  $\sigma(r) \sim 0.035$  -- This is a very direct constraint on tensors!
  - No significant evidence for  $r > 0$ . Currently  $r = 0$  and  $r = 0.1$  are at equal likelihood.
  - There may yet be a gravitational wave signal, but if there is it must be less than about half of the full signal.
- Additionally, lensing B-modes are detected at  $7.0 \sigma$  significance
- Noise in P353 is the current limiting factor and to make further progress better data at frequencies other than 150 GHz is required


# Dust Cleaned Spectrum



Dust removed points




# Results Coming soon - Keck 2014 95 GHz




For 2014 season two of the Keck array receivers switched out for 95 GHz

# Comparison of signal and noise levels




Keck 2014 95 GHz achieved noise level improves by large factor vs Planck 100 GHz

BICEP2 + Keck12+13 E-mode signal




# Keck14 95 GHz E-mode signal



Reduction in amplitude with respect to 150 GHz due to increased beam size (which is uncorrected in these map plots)



# New for 2015 - Keck220 and BICEP3



- BICEP3 receiver installed on old BICEP mount - all 95 GHz “super receiver”
- Two more Keck receivers switched out for 220 GHz
- Watch for more soon...

